

Table of Contents

Part I. Crisis Management: 1. Cardiac arrest – shockable; 2. Cardiac arrest – unshockable; 3. Paediatric ALS; 4. Intraoperative myocardial ischaemia; 5. Severe intraoperative haemorrhage; 6. Anaphylaxis; 7. Transfusion reaction; 8. Air embolism; 9. Difficult mask ventilation; 10. Unanticipated difficult airway; 11. Can't intubate, can't ventilate; 12. Laryngospasm; 13. Acutely elevated airway pressure; 14. Severe bronchospasm; 15. Aspiration; 16. Total spinal; 17. Post-partum haemorrhage; 18. Maternal collapse; 19. Newborn resuscitation; 20. Local anaesthetic toxicity; 21. Hyperkalaemia; 22. Malignant hyperthermia; 23. Terminal event checklist; Part II. Crisis Prevention: 24. 15-point machine check; 25. Elevated airway pressure; 26. Decreased oxygen saturation; 27. Increased blood pressure (hypertension); 28. Decreased blood pressure (hypotension); 29. Increased heart rate (tachycardia); 30. Decreased heart rate (bradycardia); 31. Increased CO₂ (hypercapnia); 32. Decreased CO₂ (hypocapnia); 33. Crisis prevention checklist.