

Gene–Environment Interactions for Searchers: Collaboration Between Epidemiology and Molecular Genetics

Jim van Os, Bart P.F. Rutten, and Richie Poulton

Introduction

Attempts to discover genes that relate directly to psychotic disorder (i.e., the simple “main effects” approach) have been frustrating and often disappointing, resulting in the expression of methodological concerns (Harrison and Weinberger 2005; Norton et al. 2006) (Collier 2008; Sullivan 2008; O’Donovan et al. 2008; Crow 2008). On the other hand, epidemiological research has unveiled high observed rates of schizophrenia in large cities, immigrant populations, traumatised individuals and cannabis users, at least some of which is thought to be the result of underlying environmental exposures. Exciting findings in other areas of psychiatry have motivated researchers to turn their attention to better understanding the complex ways in which nature interacts with nurture to produce psychosis. This genotype \times environmental interaction (hereafter $G \times E$) approach differs from the linear gene–phenotype approach by positing a causal role not for either genes or environment in isolation, but for their synergistic co-participation in the cause of psychosis where the effect of one is conditional on the other (EU-GEI 2008). For example, genes may moderate the psychotogenic effects of dopamine agonist drugs of abuse, or the environment may moderate the level of expression of a gene that is on the causal pathway to psychotic disorder. $G \times E$ seems a particularly suitable approach for understanding the development of psychosis because this phenotype is known to be associated with environmentally mediated risks (Cannon and Clarke 2005; Van Os et al. 2005), yet people display considerable heterogeneity in their response to those environmental exposures.

The structure of this article is as follows. First, the principles of genetic epidemiology as relevant for the study of gene–environment interaction will be reviewed briefly. Second, a brief overview will be given on what “the environment” may consist of in studies of $G \times E$ and how environmental mechanisms may be

J. van Os (✉)

Department of Psychiatry and Neuropsychology, School of Mental Health and Neuroscience, Maastricht University Medical Centre, EURON, SEARCH, Maastricht, 6200 MD, The Netherlands; Division of Psychological Medicine, Institute of Psychiatry, London SE5 8AF, UK
e-mail: j.vanos@sp.unimaas.nl

uncovered using “functional enviromics”. Third, the main $G \times E$ findings with regard to psychotic disorders will be reviewed, with a particular focus on epidemiological studies that used indirect measures of genetic risk including twin and adoption studies, family studies and psychometric risk studies. Most of the findings using direct molecular genetic measures of genetic risk will be reviewed elsewhere in this issue. Fourth, considerations will be given to possible underlying mechanisms followed by a discussion of future research and directions.

Ecogenetics

Traditional epidemiology was concerned mainly with environmental risks. Conversely, genetic researchers of complex disorders have mostly focused on molecular genetic approaches, in which the environment and interaction between genes and environment were treated as a power-reducing nuisance term. Awareness has been growing, however, that direct or indirect measures of genetic variation can be considered as a conventional epidemiological risk factor in association studies (Sham 1996) and that epidemiological theory can be readily applied to genetically sensitive data sets (Susser and Susser 1989; Ottman 1990). Thus, epidemiologists and human geneticists have been gradually integrating their respective fields of research into a new discipline called genetic epidemiology (Khoury et al. 1993). Within genetic epidemiology, the term *ecogenetics* refers to the study of specific gene–environment relationships (Motulsky 1977). Within an ecogenetic framework, several types of gene–environment relationships are relevant for the study of complex disorders, representing different biologically plausible mechanisms by which genes and environment can co-influence disease outcome (Khoury et al. 1993; Kendler and Eaves 1986; Ottman 1996; Plomin et al. 1977; Van Os and Marcelis 1998).

Ecogenetics in Psychiatry

Until recently, the conventional wisdom within psychiatry and behavioural genetics was that $G \times E$ was exceedingly rare and difficult to demonstrate. The revival of interest in $G \times E$ derives largely from (1) failures of direct gene–phenotype association studies to uncover genes related to susceptibility for psychiatric disorders and the realisation that their multifactorial aetiology likely includes many complicated interactive effects requiring more advanced approaches (Hamer 2002; Rutter 2006); (2) work demonstrating the operation of $G \times E$ in many other branches of medicine; and (3) recent evidence of $G \times E$ within psychiatry (Moffitt et al. 2005).

The recent $G \times E$ findings in psychiatry suggest that genes are likely to influence disorder mostly indirectly, via their impact upon physiological pathways, and work by increasing (or decreasing) the likelihood of developing a psychiatric disorder, rather than as direct *causes* of disorder per se. Thus, the notion of “a gene for. . .” is misleading and diverts attention from more important issues (Kendler 2005, 2006). Further, some theorists now suggest that (1) additive, non-interactive genetic effects may be less common than previously assumed (cf. Colhoun et al. 2003); (2) studying genes in isolation from known environmental risks may fail to detect impor-

tant genetic influences; and (3) traditional notions of multiplicative interaction are probably not appropriate for “real-world” interactions (Darroch 1997), particularly given the ubiquity of some environmental exposures (Moffitt et al. 2005; Rutter et al. 2006). Thus, biological synergism (co-participation of causes to some outcome) between environmental exposure and background genetic vulnerability is thought to be common in multifactorial disorders such as psychosis. The classic problem, however, is how co-participation between causes in nature (biological synergism) can be inferred from statistical manipulations with research data (statistical interaction), in particular with regard to the choice of additive (change in risk occurs by adding a quantity) or multiplicative (change in risk occurs by multiplying with a quantity) models. It has been shown that the true degree of biological synergism can be better estimated from—but is not the same as—the additive statistical interaction rather than the much more often used multiplicative interaction (Darroch 1997).

Genetic Moderation of Sensitivity to Environment

According to the concept of genetic moderation of sensitivity to the environment, differences in genetic endowment explain why people respond differently to the same environment (Fig. 1). Most evidence for this type of $G \times E$ in psychosis has come indirectly from twin and adoption studies, and a variety of naturalistic designs in which non-specific genetic contributions have been assessed. More recently, researchers have obtained information about how variation in specific measured genes interacts with specific measured environments (Moffitt et al. 2005). Genetic moderation of environmental sensitivity gives rise to *synergism*, or *interaction*, as the biological effects of G and E are dependent on each other in such a way that exposure to neither or either one alone does not result in the outcome in question, whereas exposure to both does. For example, a well-known example of gene–environment interaction is the observation that among Orientals, alcohol sensitivity is strongly regulated by genetic polymorphism of the aldehyde dehydrogenase (*ALDH2*) gene. Similarly, there is strong evidence that some polymorphisms may be involved in psychiatric disorders. For example, the gene encoding the serotonin transporter (5-HTT) contains a regulatory variation (5-HTTLPR), the short (“s”)




Fig. 1 Gene \times environment interaction: genes controlling environmental sensitivity

allele of which is associated with lower transcriptional efficiency of the promoter as compared to the long (“l”) allele. Data from animal and human research indicate that 5-HTTLPR may interact with environmental adversity to cause depression, reflecting underlying developmental mechanisms that affect the structural connectivity, and, as a consequence, functional interactions, within a neural circuit involved in the regulation of emotional reactivity and extinction of fear (Canli and Lesch 2007; Champoux et al. 2002; Wellman et al. 2007; Caspi et al. 2003; Jacobs et al. 2006) (Fig. 2).




Fig. 2 Promoter activity of the 5-HTT gene is modified by sequence elements within the proximal regulatory region; the short (“s”) allele is associated with lower transcriptional efficiency of the promoter as compared to the long (“l”) allele: converging evidence of $G \times E$ in the depression from animal to human, human to animal, and preclinical to clinical and clinical to preclinical

Although gene–environment synergism is likely prevalent, other models of disease causation, including models that imply that there is no synergism (synergism is zero), may also likely apply, although likely to a lesser degree. For example, an individual may get schizophrenia only if in possession of a certain type of vulnerability conferred by either genetic or environmental factors. An environmental factor could disrupt early brain development in the same manner as a genetic mutation. In this model, synergism is zero and the effect of genes and environment is said to be *additive*.

Environmental Impact on DNA Sequence and Methylation

Apart from genes impacting on sensitivity for environmental risk factors, $G \times E$ in psychotic disorder may also take the form of environmental factors impacting either the DNA sequence (causing de novo mutations) or DNA methylation (caus-

ing altered gene expression through epimutations). The most suggestive epidemiological evidence for such mechanisms in psychosis comes from studies linking advanced paternal age to the risk of schizophrenia in the offspring (Malaspina et al. 2001; Zammit et al. 2003; Byrne et al. 2003; Sipos et al. 2004). Paternal age varies as a function of the sociocultural environment (Weisfeld and Weisfeld 2002). The observed paternal age effect on schizophrenia may consist of mutagenesis, causing de novo spontaneous mutations, which would then propagate and accumulate in successive generations of sperm-producing cells. Alternatively, the mechanism underlying the paternal age effect may be genomic imprinting (Flint 1992). Genomic imprinting is the phenomenon whereby a small subset of all the genes in the genome is expressed according to their parent of origin. Some imprinted genes are expressed from a maternally inherited chromosome and silenced on the paternal chromosome, whereas other imprinted genes show the opposite expression pattern and are only expressed from a paternally inherited chromosome (Wilkinson et al. 2007). One of the mechanisms for gene silencing is DNA methylation. The inherited methylation pattern is maintained in somatic cells but is erased and re-established late in spermatogenesis for paternally imprinted genes, a process that could become impaired as age advances.

Although research on DNA methylation as an “epigenetic” mechanism underlying $G \times E$ in psychiatry is in an early phase, this field appears promising. For example, early maternal behaviour in animals can affect offspring stress sensitivity through altered DNA methylation of key neuronal receptor genes involved in the stress response (Weaver et al. 2004; Meaney and Szyf 2005). Environmentally induced epigenetic mechanisms may explain a range of epidemiological findings including typical age-of-onset incidence curves, monozygotic twin discordance, sex differences, possible risk-increasing effects of prenatal factors associated with in utero folate deficiency (a key component of DNA methylation) (Susser et al. 1996; Zammit et al. 2007; Smits et al. 2004) and possible risk-increasing effects of developmental trauma (Read et al. 2005). A fascinating report from Denmark is suggestive of epigenetic effects involving urban birth and upbringing. Thus, the authors demonstrated that the risk-increasing effect associated with urban birth of the older sibling “carries over” to increase the risk of schizophrenia in the next sibling who was born in a rural area (Pedersen and Mortensen 2006). This evidence is compatible with transmission of a germline epimutation associated with the urban environment. For further details on epigenetics in the context of $G \times E$, refer to the article by Oh et al. (this issue).

Gene–Environment Correlation

In contrast to $G \times E$, gene–environment correlation (hereafter rGE) refers to how differences in an individual’s genotype can “drive” differential environmental exposure (Fig. 3). In rGE , exposure to environmental events is not a random phenomenon but rather stems (at least partly) from differences in genetic make-up




Fig. 3 Gene \times environment correlation: genes controlling environmental exposure

(Plomin et al. 1977). rGEs come in three main forms. *Passive* rGE refers to environmental influences linked to genetic effects external to the person. For example, parents create the early child-rearing environment as well as provide genetic material to their offspring. Passive rGE occurs when parental behaviour, which is partly under genetic control, influences the nature of the early child-rearing environment. Thus, parental genes can exert an influence on the child via the environment, but whose effects are independent of the child itself. In contrast, *Active* rGE (e.g., selection of specific environments or “niche picking”) and *Evocative* rGE arise largely as a result of genetic factors nested within the individual (Rutter et al. 2006). Evocative rGE refers to the impact of the child’s behaviour on their social environment, in particular the responses they elicit from people around them. One person’s preference for sporting activities over another person’s penchant for artistic endeavours, thus selecting themselves into different environments, is an example of active rGE, while the different responses elicited from the social environment by gregarious versus shy individuals exemplify evocative rGE. Combining examples of rGE and $G \times E$ in one illustrative situation: rGE might manifest as arguments and disagreements preceding marital dissolution, yet $G \times E$ may determine who becomes depressed as a result of that relationship breakdown.

Confounding of $G \times E$ by rGE

In studies aimed at detecting gene–environment interactions, rGE is noise and must be ruled out. In other words, the “E” in $G \times E$ must be shown to be a true environmentally mediated effect rather than a genetic epiphenomenon. For example, does the genetic liability for schizophrenia increase the psychotogenic effect of cannabis or does schizophrenia genetic liability increase the likelihood of *using* cannabis? Experimental paradigms (see below) are able to deal effectively with this problem by randomly assigning participants to the exposed and unexposed conditions. In observational designs, however, confounding by rGE is difficult to rule out but can be tested separately. An interesting example concerns urbanicity and schizophrenia. As discussed below, four independent studies have suggested that the urban environ-

ment may contribute to the onset of psychotic disorder in individuals at genetic risk (i.e., evidence for $G \times E$). An alternative explanation, however, is that the genetic liability for schizophrenia increases the likelihood of moving to the big city, i.e., there may be rGE. A priori this is unlikely, given the fact that the effect of urbanicity on schizophrenia is restricted to the window of childhood and adolescence: children do not make the family decision to move to the big city, regardless of whether they are genetically inclined to do so or not. Two twin studies from Australia and the Netherlands on urban mobility support this notion (Whitfield et al. 2005; Willemsen et al. 2005). The Australian study showed more evidence for influence of genetic factors on urban mobility than the Dutch study. However, genetic influence in the Australian study was mostly apparent in older individuals who were well past the age at risk for onset of schizophrenia; environmental factors accounted for most of the variation in younger individuals. The reason for the discrepancy in genetic contribution to urban mobility between the Australian and the Dutch study is likely related to contextual factors. Just as the heritability of alcoholism has been shown to differ as a function of societal availability (severe restriction resulting in alcohol use only by those who are genetically most predisposed), so was the genetic influence on urban mobility shown to vary as a function of base rate of the urban outcome, which was only 10% in Australia versus around 30–50% (very heavy and heavy urbanisation) in the Netherlands. More evidence of genetic influence in Australia therefore may, in part, be the result of the lower base rate of urbanicity. Thus, the conclusion from the Australian and Dutch twin studies is that there are likely only very few human characteristics beyond any genetic influence, including urban mobility. However, in young adulthood, the age range during which psychotic disorder typically declares itself, environmental more than genetic factors may influence exposure to the risk environment that urbanicity represents (van Os 2005), making rGE unlikely.

Another important issue in rGE is that genetic effects on the outcome can be direct or indirect (Fig. 4). For example, genes may have an effect on both the outcome and the environmental exposure, while the environment has no effect on the outcome. In this case, the observed association between the environment and the outcome is genetically confounded (Fig. 4a). On the other hand, genes may have an effect on the environment, but no direct effect on the outcome, as only the environment has a causal effect (Fig. 4b). This is the situation where the environment is on the causal pathway between genes and environment, a situation that can help in providing evidence for a true causal contribution of an environmental factor to disease (Katan 1986) (referred to sometimes as “Mendelian randomisation”; Davey Smith and Ebrahim 2005). For example, evidence in the situation of Fig. 4b of an association between the gene and the outcome can only be explained if there is a true causal relationship between the environmental risk factor and the outcome. Given random assortment of genes from parents to offspring during gamete formation and conception, gene–outcome associations representing gene–causal exposure associations are not generally susceptible to the reverse causation or confounding that may plague conventional observational studies.




Fig. 4 Gene \times environment correlation (rGE): causality of environment

The Environment, Experimental Ecogenetics and Functional Enviromics

The Environment and Psychosis

Here, we refer to the environment broadly as all non-genetic influences that are associated with at least two exposure states. Sometimes, a distinction is made between “biological” and “social” environmental exposures but such a distinction may not be helpful as long as the underlying mechanisms, which are likely overlapping, are not elucidated. There are a number of environmental exposures that are associated with psychotic disorders and symptoms and for which a mechanism of gene–environment interaction has been proposed. These environmental exposures are summarised in Box 1, together with an indication to what degree the evidence for an association with schizophrenia is supported by meta-analytic estimates from systematic reviews. The most solid evidence for an association with schizophrenia and related psychosis outcomes is for paternal age, migration, urbanicity and cannabis use, the latter two particularly in the case of exposure during development.

Environmental Measurement and Experimental Ecogenetics

There are legitimate concerns about how to accurately capture the environmental risk exposure history of participants. This task is particularly challenging when measuring psychosocial risk factors whose negative effects may act cumulatively across long periods of the life course. Equally challenging are the inherent difficulties in precisely measuring “unit exposure” for illicit substances such as cannabis, which can be ingested in different forms, with different THC levels, using different methods. Measuring tobacco intake is comparatively straightforward but even this presents problems with accuracy of recall over long periods.

Box 1 Published Environmental Exposures for Psychosis for which G × E has been Suggested (M+, at least one positive meta-analytic estimate; M+/-, inconclusive meta-analytic estimate; M-, no meta-analytic estimate available)

Environmental variables with likely impact in foetal life:

1. M+: Maternal pregnancy complications, in particular foetal hypoxia and proxies for foetal folate deficiency.
2. M+/-: Prenatal maternal infection, prenatal maternal stress, prenatal maternal folate deficiency.
3. M+: Paternal age.
4. M-: Prenatal exposure to chemical agents (e.g., lead).

Environmental variables with likely impact in early life:

5. M-: Quality of early rearing environment (institutional care, school, parents).
6. M+/-: Childhood trauma (abuse or neglect).

Environmental variables with likely impact in middle childhood/adolescence:

7. M+: Urban environment during development: a variable indicating the level of population density, or size of a city within a country, of the place where the individual was growing up (between the ages of 5 and 15 years).
8. M+: Cannabis use.
9. M+: Migration.
10. M+/-: Stressful life events.
11. M-: Traumatic brain injury.

Measures of the wider social environment:

12. M-: Neighbourhood measures of social fragmentation, social capital and social deprivation.

Measures of the micro-environment in the flow of daily life:

13. M-: Small daily life stressors, assessed using momentary assessment technology, subtly impacting on affect, salience and reward.

Henquet et al. (2006) have introduced the term “experimental ecogenetics” in human psychosis research to refer to some obvious advantages: (i) randomisation precludes confounding by not only known but, critically, also unknown confounders; (ii) rGE is not an issue if “G” is randomly allocated to “E”; and (iii) it is relatively easy to make the sample size match the required power. In Fig. 5, an example is given of how the association between migration and schizophrenia, and possible genetic moderation thereof, can be examined in the context of an experimental ecogenetic design, by reducing migration to an experimental exposure of “social hostility” and by reducing the psychosis outcome to an experimental outcome of “abnormal salience attribution”, and testing the association between exposure and outcome in a genetically sensitive test design. The advent of controlled experiments with virtual reality environments may similarly represent an important asset for the study of environmental exposures (Freeman et al. 2003).

A further issue is that the environment can be conceptualised at many levels that may all be relevant to behavioural phenotypes associated with schizophrenia, varying from minor stressors in the flow of daily life as assessed by momentary assessment technologies (Myin-Germeys et al. 2001) to contextual effects of the wider social environment such as neighbourhood type or ethnic density (Kirkbride et al. 2007; Boydell et al. 2001). Finally, some environmental risks such as “urbanicity” and “ethnicity” are proxies for as-yet unidentified environmental or possibly even partly genetic factors (Pedersen and Mortensen 2006; Selten et al. 2007).




Fig. 5 Development of experimental $G \times E$ approaches

Functional Enviromics

Functional enviromics, or the study of the mechanisms underlying environmental impact on the individual to increase the risk for psychopathology, is still in its infancy, with many hypotheses yet to be tested (EU-GEI 2008). These include effects of the environment on (i) developmental programming and adult functional circuits of the brain, (ii) neuroendocrine and neurotransmitter functioning, (iii) patterns of interpersonal interactions that may shape risk for later psychopathology and (iv) affective and cognitive processing (Rutter 2005). Conversely, hypotheses need to be tested about the neural mechanism by which genetic variation may increase susceptibility to environmental stressors. These mechanisms and their underlying pathophysiological pathways need to be clarified in order to develop a priori gene–environment interaction research paradigms (Caspi and Moffitt 2006) (EU-GEI 2008). For example, it has been suggested that there may be synergistic effects of genes and environment in bringing about a “sensitisation” (Featherstone et al. 2007; Tenn et al. 2005) of mesolimbic dopamine neurotransmission (Howes et al. 2004; Collip et al. 2008). This hypothesis is supported by (i) evidence quantifying the impact of stress and dopamine agonist drugs on mesolimbic dopamine release and subsequent sensitisation (Boileau et al. 2006; Arnsten and Goldman-Rakic 1998; Covington and Miczek 2001), as well as stress-dopamine agonist cross-sensitisation (Yui et al. 2000; Nikulina et al. 2004; Hamamura and Fibiger 1993); (ii) evidence indicating that genetic risk for schizophrenia is associated with underlying alterations in the dopamine system, including increased dopamine synaptic availability (Hirvonen et al. 2005), increased striatal dopamine synthesis (Huttunen et al. 2008; Meyer-Lindenberg et al. 2002) and increased dopamine reactivity to stress (Brunelin et al. 2008; Myin-Germeys et al. 2005); and (iii) human and animal evidence that effects of environmental risk factors associated with schizophrenia have lasting effects on dopamine neurotransmission including developmental trauma (Hall et al. 1999), defeat stress associated with ethnic minority group (Covington and Miczek 2001; Tidey and Miczek 1996), prenatal hypoxia (Juarez et al. 2003, 2005; Venerosi et al. 2004) and prenatal maternal immune activation (Ozawa et al. 2006) (Meyer et al. 2008).

Thus, although there is evidence to suggest that many other neurotransmitter systems can also be targeted, a case can be made, as an example of functional enviromics, for investigating genetic variation affecting dopamine neurotransmission in interaction with environmental risk factors such as stress and dopamine agonist drugs. Molecular genetic and functional genomic studies focusing on genes associated with dopamine neurotransmission suggest that this gene group may be useful for G×E studies. For example, a recent large study focusing on gene–gene interaction (epistasis) and functional effects suggested that a network of interacting dopaminergic polymorphisms may increase risk for schizophrenia (Talkowski et al. 2008). Evidence for epistasis between genes impacting dopamine signalling can be validated using a neural systems-level intermediate phenotype approach in humans. Recent work of this type, using a prefrontal function fMRI phenotype, similarly suggests epistasis between polymorphisms in genes that control dopamine

signalling (Buckholtz et al. 2007; Meyer-Lindenberg et al. 2006). More specifically, there is evidence that schizophrenia may be characterised by a combination of prefrontal cortical dysfunction and subcortical dopaminergic disinhibition (Meyer-Lindenberg et al. 2002). Research has shown that the valine-allele carriers of a functional polymorphism in the catechol-*O*-methyltransferase gene (COMT Val¹⁵⁸Met), an important enzyme regulating prefrontal dopamine turnover, predicted increased dopamine synthesis in the midbrain, suggesting that this allele may increase the risk for schizophrenia in interaction with, for example, stress and dopamine agonist drugs (Meyer-Lindenberg et al. 2005). Several studies suggest that valine-allele carriers may indeed be more sensitive to the psychotogenic effects of drugs of abuse or stress (Henquet et al. 2006; Caspi et al. 2005; Stefanis et al. 2007).

There are examples of many other avenues that may be explored in functional enviromics. Thus, a recent systematic review suggested that more than 50% of genes potentially associated with schizophrenia, particularly AKT1, BDNF, CAPON, CCKAR, CHRNA7, CNR1, COMT, DNTBP1, GAD1, GRM3, IL10, MLC1, NOTCH4, NRG1, NR4A2/NURR1, PRODH, RELN, RGS4, RTN4/NOGO and TNF, are subject to regulation by hypoxia and/or are expressed in the vasculature (Schmidt-Kastner et al. 2006). Thus, future studies of genes proposed as candidates for susceptibility to schizophrenia should include their possible regulation by physiological or pathological hypoxia during development as well as their potential role in gene–environment interactions involving events inducing hypoxia during early development (Nicodemus et al. 2008).

Epidemiological G×E Studies in Psychosis

Epidemiological Findings

Two robust epidemiological findings suggest that “genes” and “environments” operate interactively to produce schizophrenia. First, there is widespread geographic, temporal, ethnic and other demographic variation in the incidence of schizophrenia (McGrath et al. 2004; Kirkbride et al. 2006), reinforcing the aetiological role played by environmental factors. Second, there is marked variability in people’s responses to these environmental risk factors, ranging from obvious vulnerability to extreme resilience. This well-recognised heterogeneity in response points to the operation of G×E. A number of studies have examined G×E using indirect measures of genetic risk, such as being a relative, a twin or adopted away offspring of a person with schizophrenia, or the level of psychometric psychosis proneness in a person as an expression of distributed genetic risk for psychotic disorder (see below). The advantage of these studies is that the measure of genetic risk, while non-specific and therefore not able to capture gene–environment interactions with very specific mechanisms, is nevertheless (i) well validated and (ii) represents the complete net genetic load including all gene–gene interactions. While newer studies using direct molecular genetic measures of genetic risk have the advantage of using

specific measures, they are also prone to false-positive findings, given the enormous amount of molecular genetic variation that can be used for $G \times E$ modelling, and the absence of all other factors influencing genetic risk in the model of $G \times E$ using a small contribution to genetic variation in the form of an SNP. Therefore, epidemiological studies using indirect measures of genetic risk remain useful and may point the way to $G \times E$ studies using direct measures of genetic risk; to date, they remain the most informative. A review of these findings is presented here.

Findings from Twin, Adoption and Family Studies

Twin and adoption studies provide strong but non-specific evidence for the involvement of both genes and environmental factors in the aetiology of schizophrenia (Gottesman and Shields 1976). Both have shown moderate-to-high heritability for schizophrenia but even monozygotic twins show only 50% concordance, underscoring the likelihood of environmental influences and $G \times E$ synergism for producing psychotic symptoms and disorder (Van Os and Sham 2003). Findings from several adoption studies are consistent with $G \times E$ in the development of psychotic disorders. For example, Carter et al. (2002) compared, in a 25-year longitudinal study, 212 children of schizophrenic mothers with 99 children of normal parents in terms of exposure to environmental risk (i.e., institutional care and family instability). Very few cases of psychosis were identified in those families without a history of schizophrenia but, among those with a family history, strong environmental effects were observed. Consistent with this, Tienari et al. (2004) compared adopted-away offspring ($N = 145$) of mothers with a history of psychotic illness with those without illness ($N = 158$). Measures of the rearing environment in the adoptive home were obtained (measures on scales of “critical/conflictual”, “constricted” and “boundary problems”) and revealed strong effects for those with a biological predisposition (odds ratio around 10), which were absent in those with low genetic risk (odds ratio around 1).

Findings in support of $G \times E$ also come from migration designs, which, for example, have demonstrated a higher risk of psychosis among Caribbean immigrants to the United Kingdom compared to the majority population in the United Kingdom (Cantor-Graae and Selten 2005). Further, family studies of UK-born Afro-Caribbeans have demonstrated a particularly high risk of schizophrenia among the siblings of young, Afro-Caribbean patients (15.9% compared to 1.8% in siblings of white patients), whereas the rates of schizophrenia among the white and Afro-Caribbean parents were similar (8.4 and 8.9%, respectively) (Sugarman and Craufurd 1994).

Studies Using a Psychometric Psychosis Liability Approach

Subtle subclinical expression of psychosis can be measured in the general population (Van Os et al. 2000). There is evidence that this phenotype of “psychometric

psychosis proneness” represents in part the distributed genetic risk for psychotic disorder, suggesting it could be used as a proxy to represent the factor “G” in studies of G×E, although to the degree that environmental factors contribute to the psychometric psychosis proneness measure these cannot be excluded as a source of confounding. Thus, Vollema et al. (2002) Vollema and colleagues reported that scores on the positive dimension of a schizotypy questionnaire administered to relatives of patients with psychotic disorders corresponded to their genetic risk of psychosis. Fanous et al. (2001) demonstrated that interview-based positive and negative symptoms in schizophrenia predicted their equivalent subclinical symptom dimensions in non-psychotic relatives, implying an aetiological continuum between the subclinical and the clinical psychosis phenotypes. Kendler and Hewitt (1992) studied twins from the general population and concluded that the variance in most self-reported schizotypy scales, except for perceptual aberration, involved substantial genetic contributions. MacDonald et al. (2001) found in their general population-based twin study only one common schizotypy factor, mainly explained by perceptual aberration, magical ideation, schizotypal cognitions and, to a lesser extent, social anhedonia. The common schizotypy factor was influenced by shared environmental, non-shared environmental and possibly genetic effects. Recently, a general population female twin study by Linney et al. (2003) showed that additive genetic and unique environmental effects influenced self-reported psychotic experiences. The multivariate structural equation model generated two independent latent factors, namely a positive (i.e., cognitive disorganisation, unusual experiences and delusional ideation) and a negative dimension (i.e., cognitive disorganisation and introverted anhedonia), suggesting different aetiological mechanisms for the various scales of the subclinical psychosis phenotype. In a recent, general population study using both self-reported and interview-based measures of positive and negative dimensions of psychotic experiences in 257 subjects belonging to 82 families, significant family-specific variations for both positive and negative subclinical psychosis dimensions were demonstrated, with between-family proportions of total variance between 10 and 40%. Thus, both the positive and the negative dimensions of subclinical psychosis show familial clustering in samples unselected for psychiatric disease (Hanssen et al. 2006). Operationalising the genetic effect “G” along these lines, Henquet et al. (2005) showed that a psychometric measure of psychosis proneness interacted with cannabis use to predict the likelihood of developing psychotic symptoms. In this study, rGE was unlikely to have been a confounder since no association between baseline psychosis proneness and subsequent use of cannabis was observed. Nonetheless, confounding cannot be ruled out entirely because the proxy genetic measure of psychometric psychosis proneness will also be influenced by environmental factors. As a complement to the observational designs described above, Verdoux et al. (2003) used a quasi-experimental “experience sampling” method, and obtained similar findings showing that psychosis liability moderated the effect of cannabis in terms of “switching on” psychotic symptoms in the flow of daily life. For more details on possible gene × cannabis interactions, we refer to the article by Henquet et al. (this issue). Other studies using psychometric psychosis liability as a proxy measure for genetic risk were able to demonstrate G×E

with childhood urbanicity (Spauwen et al. 2006a) (see below for more details) and childhood trauma (Spauwen et al. 2006b).

Summary of Epidemiological G×E Studies to Date

In Table 1, the different epidemiological G×E studies are summarised. For each study, the proxy genetic factor, the proxy environmental factor and the main findings as well as main limitations are summarised. Environmental exposures used in G×E studies include migration, urbanicity, obstetric complications, cannabis, stress, developmental trauma and others. In most studies, the effect of genes and environment alone was rather small, and the bulk of their effect mediated through gene–environment interactions.

Epidemiological Replications of Gene–Urbanicity Interaction

The finding that the rate of psychotic disorder is higher in children and adolescents growing up in an urban environment is well replicated (Krabbendam and van Os 2005) and unlikely to be confounded entirely by rGE due to selective drift to urban areas in those at genetic risk for psychosis (Van Os 2004; Pedersen and Mortensen 2001), although rGE may operate to some degree (Pedersen and Mortensen 2006; Selten et al. 2007) as it will in the case of many environmental risks (van Os et al. 2005). “Urbanicity” is a proxy for an as-yet unidentified environmental factor(s) prevalent in urban areas and, if causal, may contribute to up to 20–30% of the incidence of psychotic disorder in some countries (Van Os 2004). For this reason, urbanicity is an interesting factor to study in the context of G×E. Four studies in the Netherlands, Germany, Israel and Denmark have attempted to examine gene–urbanicity interactions using epidemiological designs and indirect measures of genetic risk (Spauwen et al. 2006; Van Os et al. 2003, 2004; Weiser et al. 2007). All studies found evidence for gene–urbanicity interaction and are summarised in Table 2. Clearly, the possibility of interaction between an environmental exposure in urban areas and genetic risk is in need of further study, focusing on (i) the precise nature of the urban exposure, for example, growing up in an area lacking in trust and cohesion, (ii) the psychological and neurobiological mechanism of the environmental exposure in order to develop rational hypotheses about gene–environment interaction, (iii) the nature of the genetic variation involved, and ultimately (iv) the mechanism of the gene–environment interactions.

Future Prospects

To date, the study of gene–environment interactions has largely been epidemiological, where genotype, risk exposure and disorder are studied as they occur in the population (Khoury et al. 2004). A key contribution of a robust G×E comes from

Table 1 First-generation studies of proxy gene–environment interaction in psychotic disorder

Proxy genetic variable	Proxy environmental variable	Findings	Remarks
Positive family history (FH)	Ethnic group	Familial morbid risk for psychotic disorder higher in siblings of African-Caribbean probands than in those of white probands (Sugarman and Craufurd 1994; Hutchinson et al. 1996)	➤ May be informative; however, preferably environmental exposure status and clinical status are measured in both cases <i>and</i> all first-degree relatives and analyses are adjusted for age, sex and number of relatives
	Urban birth	No association between urban birth and a positive family history for psychotic disorder (Mortensen et al. 1999); however, interaction was tested on multiplicative rather than additive scale (see below)	➤ Level of misclassification may be high because many unaffected relatives may carry the high-risk genotype
	Obstetric complications	Mostly inconclusive findings with regard to family history (Nimgaonkar et al. 1988; Kunugi et al. 1996; O'Callaghan et al. 1992)	➤ Absence of an association between positive family history and environmental exposure does <i>not</i> rule out gene–environment interaction, and <i>presence</i> of an association does <i>not</i> rule out <i>lack</i> of gene–environment interaction (Marcelis et al. 1998)
	Birth in winter/spring	Positive, negative and inconclusive associations with family history (Baron and Gruen 1988; Shur 1982; Pulver et al. 1992; Dassa et al. 1996)	➤ Evidence can be considered stronger if replicated (e.g., urbanicity findings)
	Stressful life events	Positive association with family history (Van Os et al. 1994)	➤ Testing for interaction on additive scale likely more informative (Darroch 1997)
	Urbanicity	Evidence for synergism between urban environment (proxy environmental risk) and family history (proxy genetic risk) when tested on additive scale (Van Os et al. 2003)	

Table 1 (continued)

Proxy genetic variable	Proxy environmental variable	Findings	Remarks
Having an identical twin with psychotic disorder	Being discordant for psychotic disorder	Children of both affected and non-affected twin in discordant pair have higher rate of psychotic disorder (Kringlen and Cramer 1989; Fischer 1971; Gottesman and Bertelsen 1989)	➤ Suggests environmental factor is necessary for the expression of high-risk genotype in affected twin or inhibition of protective genotype in unaffected twin
Biological parent with psychotic disorder	Growing up in dysfunctional adoptive family environment	Risk of psychotic disorder spectrum disorder or psychotic disorder-associated thought disorder higher in high-risk adoptees who had been brought up in dysfunctional adoptive family environment (Tienari et al. 2004, 1994; Wahlberg et al. 1997, 2004).	➤ Risk of psychotic disorder spectrum disorder 3% in the absence of environmental risk and 62% in the presence of environmental risk. This difference seems extremely high ➤ Children destined to develop psychotic disorder may have contributed to dysfunctional family environment rather than the other way round
	Institutional care, family instability	Very few cases of psychosis were identified in those families without a history of psychotic disorder but, among those with a family history, strong environmental effects were observed (Carter et al. 2002)	➤ Case–control comparison difficult as many other factors may be involved

Table 1 (continued)

Proxy genetic variable	Proxy environmental variable	Findings	Remarks
Having neither, one or two parents with psychotic disorder	Having positive relationships with father and mother	High-risk children with positive parental relationships had lower risk for developing psychotic disorder (Carter et al. 1999)	➤ May suggest a negative G×E
	Obstetric complications	The greater the proxy genetic risk, the greater the effect of obstetric complications, in particular foetal hypoxia, on ventricular enlargement (the psychotic disorder endophenotype) (Cannon et al. 1993, 2002)	➤ Genetic risk may increase the risk of obstetric complication (rGE).
	Paternal absence	Higher rate of paternal absence in children who subsequently developed psychotic disorder (Walker 1981)	➤ Genetic risk may increase the risk of heavy alcohol consumption or head injury resulting in greater OC effect sizes ➤ Status of electrodermal abnormality as a marker of genetic risk for psychotic disorder unclear
Having a parent with psychotic disorder and additionally having an electrodermal abnormality as a child	Sharing the same chorion with the co-twin	Concordance rate was higher for MZ twins whose marker suggested they were monozygotic than those whose marker indicated they were dizygotic (Davis and Phelps 1995)	➤ These results are compatible with an environmental factor in the prenatal environment facilitating expression of genetic risk for psychotic disorder

Table 1 (continued)

Proxy genetic variable	Proxy environmental variable	Findings	Remarks
Having expression of genetically influenced psychometric psychosis liability	Early trauma	Evidence that trauma and psychometric psychosis liability synergistically increase the risk for psychosis persistence over time (Spauwen et al. 2006)	➤ Difficult to disentangle rGE from G×E ➤ Psychometric psychosis liability very indirect measure of genetic risk
	Cannabis use	Evidence that cannabis and psychometric psychosis liability synergistically increase the risk for psychosis persistence over time (Henquet et al. 2005; see also Verdoux et al. 2003)	
	Growing up in urban environment	Evidence that urbanicity and psychometric psychosis liability synergistically increase the risk for psychosis persistence over time (Spauwen et al. 2006)	

Table 1 (continued)

Proxy genetic variable	Proxy environmental variable	Findings	Remarks
Being a member of a schizophrenia pedigree	Traumatic brain injury	Within the schizophrenia pedigrees but not bipolar pedigrees, traumatic brain injury was associated with a greater risk of schizophrenia, consistent with synergistic effects between genetic vulnerability for schizophrenia and traumatic brain injury	➤ Similar comments as for positive family history
None	Having an older father	Having an older father is associated with an increased risk of schizophrenia in the offspring (Malaspina et al. 2001; Zammit et al. 2003; Byrne et al. 2003; Sipos et al. 2004)	➤ The underlying mechanism of this association may represent a special case of gene–environment interaction whereby the environment impacts on DNA sequence (de novo mutation) or DNA methylation (affecting gene expression). Thus, age of the father is a variable that is partly under control from the sociocultural environment, and older age may have an effect on DNA methylation in sex cells (the inherited methylation pattern in humans is maintained in somatic cells but is erased and re-established late in spermatogenesis for paternally imprinted genes, a process that could become impaired as age advances). Alternatively, advanced paternal age may lead to an increased rate of de novo mutations in gametes

Table 2 Studies of gene–urbanicity interactions

Study	Country	Measure genetic risk	Measure urbanicity	Psychosis outcome	Rate unexposed ^a	Rate E ^b	Rate G ^c	Rate GE ^d
Van Os et al. (2003)	The Netherlands	Family history psychosis	Population density – dichotomous	Psychotic disorder	0.85%	1.59%	3.01%	9.72%
Van Os et al. (2004)	Denmark	Family history psychotic disorder	Five categories from capital city to rural area – five levels	Psychotic disorder	Summary increase in incidence associated with urbanicity in individuals WITHOUT family history: 0.054%	Summary increase in incidence associated with urbanicity in individuals WITH family history: 0.22%	Summary increase in incidence associated with urbanicity in individuals WITH family history: 0.22%	Summary increase in incidence associated with urbanicity in individuals WITH family history: 0.22%
Spauwen et al. (2006)	Germany	Psychometric psychosis liability	City of Munich versus surrounding villages – dichotomous	Psychotic symptoms	14.2%	12.1%	14.9%	29%
Weiser et al. (2007)	Israel	Cognitive impairment endophenotype	Population density – five levels	Psychotic disorder	Summary increase in incidence associated with urbanicity in cognitively non-vulnerable group: 0.011%	Summary increase in incidence associated with urbanicity in cognitively non-vulnerable group: 0.011%	Summary increase in incidence associated with urbanicity in cognitively vulnerable group: 0.10%	Summary increase in incidence associated with urbanicity in cognitively vulnerable group: 0.10%

^aThose exposed to neither urbanicity nor genetic risk.^bThose exposed to urbanicity only.^cThose exposed to genetic risk only.^dThose exposed to both urbanicity and genetic risk.

knowing that three apparently unconnected factors (gene, environmental risk factor and disorder) are in fact causally linked (Moffitt et al. 2005). However, there are a number of methodological concerns that continue to challenge genetic-epidemiological research, mainly because observational methods struggle to achieve the degree of control that is possible using experimental designs (EU-GEI 2008; Caspi and Moffitt 2006). Concerns are listed below.

The Ideal Sample Size for G×E Research

Clearly the optimal sample size required to detect G×E will vary according to the design used. For example, case-control studies will generally require very large sample sizes simply because the genetic effects are expected to be small. However, even with prospective cohort studies, large sample sizes may be required when the environmental risk factor(s) and/or disorder of interest occur at low frequencies. However, large sample sizes are not always necessary, or desirable given the costs of amassing large samples. Indeed, sample size requirements can be substantially reduced with high-quality measurement of environmental risk factors, especially when measures are repeated over time (Wong et al. 2003); in particular, the use of momentary assessment technologies with many repeated measures holds promise for the detection of subtle gene-environment interactions (Myin-Germeys et al. 2001; Wichers et al. 2007a, b). Other methods to reduce sample size, based on selection of extreme exposure groups, may also apply (Boks et al. 2007).

Biostatistics

It is likely that mass genome-wide molecular genetic approaches, “enriched” with a few measures of “environmental” exposures, will create invalid and confusing findings, largely because of the extent of multiple testing and the opportunities for post hoc analyses afforded by such studies. It is of paramount importance to consider the study of G×E as a separate discipline, requiring a highly specialised and multidisciplinary approach taking both environment and genes seriously. A hypothesis-driven strategy focusing on final common pathways in which biological synergism between genetic and environmental mechanisms takes place, fed by information from functional enviromics and functional genomics pointing to promising neural systems and processes, may constitute the most productive approach. In combination, this will enable a translational approach for systematically studying the effect of environmental manipulations on neural systems linked to genetic risk for schizophrenia. However, even a hypothesis-driven approach is likely to face major challenges in the area of biostatistics. Even allowing for, as discussed earlier, the major problem of how to bridge the gap between statistical interaction (statistical manipulations of data) and biological synergism (biological processes in nature), which currently cannot be estimated directly (Van Os and Sham 2003), solutions to, for example,

modelling multiple ambiguous haplotype \times environment interactions need to be developed (Lake et al. 2003). Fortunately, software allowing for modelling complicated interactions is currently being incorporated in several statistical programs (Li and Stephens 2003; Lange et al. 2004).

Which Endophenotypes to Study?

In order to elucidate converging pathways that are the site of biological synergism between genes and environments, a wide range of approaches employing intermediate (or endo-) phenotypes may be used. For example, one may focus on the domain of neural systems-level intermediate phenotypes (Meyer-Lindenberg et al. 2006; Murray et al. 2008; Barkus et al. 2007), cognition (Filbey et al. 2008; Touloupoulou et al. 2007; Barnett et al. 2007a, b; Bombin et al. 2008), neuroanatomy (van Haren et al. 2008; Boos et al. 2007; Marcelis et al. 2003), salience attribution (Jensen et al. 2008; Kapur 2003), treatment response (Arranz and de Leon 2007), measures of course and outcome (Verdoux et al. 1996), subclinical psychosis expression (Stefanis et al. 2004; Schurhoff et al. 2003, 2007), neurotic symptoms (Zinkstok et al. 2008) and dynamic cerebral phenotypes in early-onset groups (Arango et al. 2008). The appeal of studying endophenotypes is obvious in that, compared to clinical diagnoses which are often characterised by substantial heterogeneity, endophenotypes appear to be cleaner, simpler constituents of psychopathology and (maybe falsely) promise improved chances of detecting true gene effects. Nonetheless, questions remain about which endophenotypes, for which disorder, are most worthy of study in a $G \times E$ framework. One argument against the use of endophenotypes is their apparently lower heritability estimates than the clinical phenotype (Greenwood et al. 2007). Although at first glance this may seem a valid argument, lower heritability estimates are only to be expected if endophenotypes reflect the “pure” contribution of genes and the clinical phenotype additionally represents the contribution of gene–environment interactions. The reason for this is that heritability estimates are derived from genetic epidemiological studies that estimate simple genetic and simple environmental contributions to schizophrenia liability. Unfortunately, these studies do not model the contribution of gene–environment interactions ($G \times E$), because researchers tend to not include direct measures of the environment in such studies, thus precluding the quantification of gene–environment interactions. Therefore, the heritability of schizophrenia may be 80%, but simulations show that gene–environment interactions may make up the bulk of this proportion (Van Os and Sham 2003). Thus, endophenotypes may be more suitable measures of “pure” genetic risk, as heritability estimates of the clinical disorder may be inflated by gene–environment interactions. Further research on this issue is needed.

Multiple Tests

As mentioned earlier, there are legitimate concerns about low prior probability testing for associations between a large number of polymorphisms (for example, via

SNP chips) and specific disorders in the absence of some guiding theory that will allow researchers to sort true from false-positive associations. Guarding against “fishing trips” is important if we are to advance our understanding of how G×E operates in the development of schizophrenia.

Conclusion

Not only is there meta-analytic support for environmental effects on schizophrenia risk, evidence is now accumulating that environmental exposures are impacting on the risk for psychotic disorder in co-participation with genetic factors and that effects of genes and environment in isolation are likely small or non-existent.

Embracing a G×E approach has implications for gene discovery. That is, selecting and/or stratifying samples based on documented environmental risk exposure may help not only in the quest to identify new susceptibility genes for psychotic disorders but also in unravelling the pathway(s) to the onset of first-episode psychosis. For molecular genetic research, this means that the strategy of “brute force” (Collier 2008), used to compensate for loss of power due to underlying G×E by inclusion of huge samples of many thousands of patients and hundreds of thousands of markers along the genome, may be complemented by imaginative approaches based on environmental stratification. Genetic odds ratio of 1.1 in non-stratified samples may be considerably higher in exposed samples. In addition, distal tiny genetic contributions by themselves explain little if more proximal interactions with environmental component causes, explaining the underlying pathophysiology.

It is obvious that more funding needs to be directed to G×E research – after nearly 1,500 inconclusive molecular genetic investigations in schizophrenia complementary approaches no longer need to be excluded. The European Network of Schizophrenia Networks for the Study of Gene–Environment Interactions (EU-GEI) (EU-GEI 2008) has suggested that part of the funding may be necessary to bring together the multitude of disciplines, currently working in isolation of each other, which is necessary for the study of gene–environment interactions.

Future research needs to better integrate epidemiological and experimental paradigms focusing on functional enviromics and functional genomics (Caspi and Moffitt 2006; EU-GEI 2008). This is desirable because neither traditional genetic epidemiology nor epidemiologic studies on isolated environmental factors can tell us much about the biological mechanisms involved in a G×E. These approaches are complementary, with each informing the other, and ideally should be used in unison for best effect. Many (but by no means all) of the challenges confronting genetic epidemiology listed above can be addressed using experimental designs with their advantages of greater experimental control and precision. However, these benefits have to be balanced against the loss of ecological validity that can sometimes result.

Epidemiologists should be encouraged to incorporate more physiological (i.e., mechanistic) measures in their studies and to move beyond two-way interactions to models involving multiple genes and environments, as well as gene–gene and environment–environment interactions.

Acknowledgment Parts of this chapter appeared as: van Os, J., Poulton, R. (2008). Environmental vulnerability and genetic-environmental interactions. In: *The Recognition and Management of Early Psychosis: A Preventive Approach*, 2nd edition (eds. H. Jackson and P. McGorry). Cambridge University Press: Cambridge; van Os J, Rutten BP, Poulton R. Gene-environment interactions in schizophrenia: review of epidemiological findings and future directions. *Schizophrenia Bulletin* 34:1066–1082, 2009.

References

- Arango C, Moreno C, Martinez S, Parellada M, Desco M, Moreno D, Fraguas D, Gogtay N, James A, Rapoport J. (2008). Longitudinal brain changes in early-onset psychosis. *Schizophrenia Bulletin* 34:341–53.
- Arnsten AF, Goldman-Rakic PS. (1998). Noise stress impairs prefrontal cortical cognitive function in monkeys: evidence for a hyperdopaminergic mechanism. *Arch Gen Psychiatry* 55:362–8.
- Arranz MJ, de Leon J. (2007). Pharmacogenetics and pharmacogenomics of schizophrenia: a review of last decade of research. *Mol Psychiatry* 12:707–47.
- Barkus E, Stirling J, Hopkins R, McKie S, Lewis S. (2007). Cognitive and neural processes in non-clinical auditory hallucinations. *Br J Psychiatry Suppl* 51:s76–s81.
- Barnett JH, Heron J, Ring SM, Golding J, Goldman D, Xu K, Jones PB. (2007b). Gender-specific effects of the catechol-*O*-methyltransferase Val108/158Met polymorphism on cognitive function in children. *Am J Psychiatry* 164:142–9.
- Barnett JH, Jones PB, Robbins TW, Muller U. (2007a). Effects of the catechol-*O*-methyltransferase Val158Met polymorphism on executive function: a meta-analysis of the Wisconsin Card Sort Test in schizophrenia and healthy controls. *Mol Psychiatry* 12:502–9.
- Baron M, Gruen R. (1988). Risk factors in schizophrenia. Season of birth and family history [see comments]. *Br J Psychiatry* 152:460–5.
- Boileau I, Dagher A, Leyton M, Gunn RN, Baker GB, Diksic M, Benkelfat C. (2006). Modeling sensitization to stimulants in humans: an [¹¹C]raclopride/positron emission tomography study in healthy men. *Arch Gen Psychiatry* 63:1386–95.
- Boks MP, Schipper M, Schubart CD, Sommer IE, Kahn RS, Ophoff RA. (2007). Investigating gene environment interaction in complex diseases: increasing power by selective sampling for environmental exposure. *Int J Epidemiol* 36:1363–9.
- Bombin I, Arango C, Mayoral M, Castro-Fornieles J, Gonzalez-Pinto A, Gonzalez-Gomez C, Moreno D, Parellada M, Baeza I, Graell M, Otero S, Saiz PA, Patino-Garcia A. (2008). DRD3, but not COMT or DRD2, genotype affects executive functions in healthy and first-episode psychosis adolescents. *Am J Med Genet B Neuropsychiatr Genet* 147:873–9.
- Boos HB, Aleman A, Cahn W, Pol HH, Kahn RS. (2007). Brain volumes in relatives of patients with schizophrenia: a meta-analysis. *Arch Gen Psychiatry* 64:297–304.
- Boydell J, van Os J, McKenzie K, Allardyce J, Goel R, McCreadie RG, Murray RM. (2001). Incidence of schizophrenia in ethnic minorities in London: ecological study into interactions with environment. *BMJ* 323:1336–8.
- Brunelin J, d'Amato T, van Os J, Cochet A, Suaud-Chagny MF, Saoud M. (2008). Effects of acute metabolic stress on the dopaminergic and pituitary–adrenal axis activity in patients with schizophrenia, their unaffected siblings and controls. *Schizophr Res* 100:206–211.
- Buckholtz JW, Sust S, Tan HY, Mattay VS, Straub RE, Meyer-Lindenberg A, Weinberger DR, Callicott JH. (2007). fMRI evidence for functional epistasis between COMT and RGS4. *Mol Psychiatry* 12:893–5, 885.
- Byrne M, Agerbo E, Ewald H, Eaton WW, Mortensen PB. (2003). Parental age and risk of schizophrenia: a case–control study. *Arch Gen Psychiatry* 60:673–8.
- Canli T, Lesch KP. (2007). Long story short: the serotonin transporter in emotion regulation and social cognition. *Nat Neurosci* 10:1103–9.

- Cannon M, Clarke MC. (2005). Risk for schizophrenia – broadening the concepts, pushing back the boundaries. *Schizophr Res* 79:5–13.
- Cannon TD, Mednick SA, Parnas J, Schulsinger F, Praestholm J, Vestergaard A. (1993). Developmental brain abnormalities in the offspring of schizophrenic mothers. I. Contributions of genetic and perinatal factors [see comments]. *Arch Gen Psychiatry* 50:551–64.
- Cannon TD, van Erp TG, Rosso IM, Huttunen M, Lonnqvist J, Pirkola T, Salonen O, Valanne L, Poutanen VP, Standertskjold-Nordenstam CG. (2002). Fetal hypoxia and structural brain abnormalities in schizophrenic patients, their siblings, and controls. *Arch Gen Psychiatry* 59:35–41.
- Cantor-Graae E, Selten JP. (2005). Schizophrenia and migration: a meta-analysis and review. *Am J Psychiatry* 162:12–24.
- Carter JW, Parnas J, Cannon TD, Schulsinger F, Mednick SA. (1999). MMPI variables predictive of schizophrenia in the Copenhagen High-Risk Project: a 25-year follow-up. *Acta Psychiatr Scand* 99:432–40.
- Carter JW, Schulsinger F, Parnas J, Cannon T, Mednick SA. (2002). A multivariate prediction model of schizophrenia. *Schizophr Bull* 28:649–82.
- Caspi A, Moffitt TE, Cannon M, McClay J, Murray R, Harrington H, Taylor A, Arseneault L, Williams B, Braithwaite A, Poulton R, Craig IW. (2005). Moderation of the effect of adolescent-onset cannabis use on adult psychosis by a functional polymorphism in the catechol-O-methyltransferase gene: longitudinal evidence of a gene \times environment interaction. *Biol Psychiatry* 57:1117–27.
- Caspi A, Moffitt TE. (2006). Gene–environment interactions in psychiatry: joining forces with neuroscience. *Nat Rev Neurosci* 7:583–90.
- Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, McClay J, Mill J, Martin J, Braithwaite A, Poulton R. (2003). Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* 301:386–9.
- Champoux M, Bennett A, Shannon C, Higley JD, Lesch KP, Suomi SJ. (2002). Serotonin transporter gene polymorphism, differential early rearing, and behavior in rhesus monkey neonates. *Mol Psychiatry* 7:1058–63.
- Colhoun HM, McKeigue PM, Davey Smith G. (2003). Problems of reporting genetic associations with complex outcomes. *Lancet* 361:865–72.
- Collier DA. (2008). Schizophrenia: the polygene princess and the pea. *Psychol Med* 1–5.
- Collip D, Myin-Germeys I, Van Os J. (2008). Does the concept of “sensitization” provide a plausible mechanism for the putative link between the environment and schizophrenia? *Schizophr Bull* 34:220–5.
- Covington HE, 3rd, Miczek KA. (2001). Repeated social-defeat stress, cocaine or morphine. Effects on behavioral sensitization and intravenous cocaine self-administration “binges”. *Psychopharmacology (Berl)* 158:388–98.
- Crow TJ. (2008). The emperors of the schizophrenia polygene have no clothes. *Psychol Med* 1–5.
- Darroch J. (1997). Biologic synergism and parallelism [see comments]. *Am J Epidemiol* 145:661–8.
- Dassa D, Sham PC, Van Os J, Abel K, Jones P, Murray RM. (1996). Relationship of birth season to clinical features, family history, and obstetric complication in schizophrenia. *Psychiatry Res* 64:11–7.
- Davey Smith G, Ebrahim S. (2005). What can mendelian randomisation tell us about modifiable behavioural and environmental exposures? *BMJ* 330:1076–9.
- Davis JO, Phelps JA. (1995). Twins with schizophrenia: genes or germs? *Schizophr Bull* 21:13–8.
- EU-GEI. (2008). European Network of Schizophrenia Networks for the Study of Gene Environment Interactions. Schizophrenia aetiology: do gene–environment interactions hold the key? *Schizophr Res* 102:21–6.
- Fanous A, Gardner C, Walsh D, Kendler KS. (2001). Relationship between positive and negative symptoms of schizophrenia and schizotypal symptoms in nonpsychotic relatives. *Arch Gen Psychiatry* 58:669–673.

- Featherstone RE, Kapur S, Fletcher PJ. (2007). The amphetamine-induced sensitized state as a model of schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 31:1556–71.
- Filbey FM, Touloupoulou T, Morris RG, McDonald C, Bramon E, Walshe M, Murray RM. (2008). Selective attention deficits reflect increased genetic vulnerability to schizophrenia. *Schizophr Res* 101:169–75.
- Fischer M. (1971). Psychoses in the offspring of schizophrenic monozygotic twins and their normal co-twins. *Br J Psychiatry* 118:43–52.
- Flint J. (1992). Implications of genomic imprinting for psychiatric genetics. *Psychol Med* 22:5–10.
- Freeman D, Slater M, Bebbington PE, Garety PA, Kuipers E, Fowler D, Met A, Read CM, Jordan J, Vinayagamoorthy V. (2003). Can virtual reality be used to investigate persecutory ideation? *J Nerv Ment Dis* 191:509–14.
- Gottesman, II, Bertelsen A. (1989). Confirming unexpressed genotypes for schizophrenia. Risks in the offspring of Fischer's Danish identical and fraternal discordant twins [see comments]. *Arch Gen Psychiatry* 46:867–72.
- Gottesman, II, Shields J. (1976). A critical review of recent adoption, twin, and family studies of schizophrenia: behavioral genetics perspectives. *Schizophr Bull* 2:360–401.
- Greenwood TA, Braff DL, Light GA, Cadenhead KS, Calkins ME, Dobie DJ, Freedman R, Green MF, Gur RE, Gur RC, Mintz J, Nuechterlein KH, Olincy A, Radant AD, Seidman LJ, Siever LJ, Silverman JM, Stone WS, Swerdlow NR, Tsuang DW, Tsuang MT, Turetsky BI, Schork NJ. (2007). Initial heritability analyses of endophenotypic measures for schizophrenia: the consortium on the genetics of schizophrenia. *Arch Gen Psychiatry* 64:1242–50.
- Hall FS, Wilkinson LS, Humby T, Robbins TW. (1999). Maternal deprivation of neonatal rats produces enduring changes in dopamine function. *Synapse* 32:37–43.
- Hamamura T, Fibiger HC. (1993). Enhanced stress-induced dopamine release in the prefrontal cortex of amphetamine-sensitized rats. *Eur J Pharmacol* 237:65–71.
- Hamer D. (2002). Genetics. Rethinking behavior genetics. *Science* 298:71–2.
- Hanssen M, Krabbendam L, Vollema M, Delespaul P, Van Os J. (2006). Evidence for instrument and family-specific variation of subclinical psychosis dimensions in the general population. *J Abnorm Psychol* 115:5–14.
- Harrison PJ, Weinberger DR. (2005). Schizophrenia genes, gene expression, and neuropathology: on the matter of their convergence. *Mol Psychiatry* 10:40–68; image 5.
- Henquet C, Krabbendam L, Spauwen J, Kaplan C, Lieb R, Wittchen HU, van Os J. (2005). Prospective cohort study of cannabis use, predisposition for psychosis, and psychotic symptoms in young people. *BMJ* 330:11.
- Henquet C, Rosa A, Krabbendam L, Papiol S, Fananas L, Drukker M, Ramaekers JG, van Os J. (2006). An experimental study of catechol-o-methyltransferase Val158Met moderation of delta-9-tetrahydrocannabinol-induced effects on psychosis and cognition. *Neuropsychopharmacology* 31:2748–57.
- Hirvonen J, van Erp TG, Huttunen J, Aalto S, Nagren K, Huttunen M, Lonnqvist J, Kaprio J, Hietala J, Cannon TD. (2005). Increased caudate dopamine D2 receptor availability as a genetic marker for schizophrenia. *Arch Gen Psychiatry* 62:371–8.
- Howes OD, McDonald C, Cannon M, Arseneault L, Boydell J, Murray RM. (2004). Pathways to schizophrenia: the impact of environmental factors. *Int J Neuropsychopharmacol* 7(Suppl 1):S7–S13.
- Hutchinson G, Takei N, Fahy TA, Bhugra D, Gilvarry C, Moran P, Mallett R, Sham P, Leff J, Murray RM. (1996). Morbid risk of schizophrenia in first-degree relatives of white and African-Caribbean patients with psychosis. *Br J Psychiatry* 169:776–80.
- Huttunen J, Heinimaa M, Svirskis T, Nyman M, Kajander J, Forsback S, Solin O, Ilonen T, Korkeila J, Ristkari T, McGlashan T, Salokangas RK, Hietala J. (2008). Striatal dopamine synthesis in first-degree relatives of patients with schizophrenia. *Biol Psychiatry* 63:114–7.
- Jacobs N, Kenis G, Peeters F, Derom C, Vlietinck R, van Os J. (2006). Stress-related negative affectivity and genetically altered serotonin transporter function: evidence of synergism in shaping risk of depression. *Arch Gen Psychiatry* 63:989–96.

- Jensen J, Willeit M, Zipursky RB, Savina I, Smith AJ, Menon M, Crawley AP, Kapur S. (2008). The formation of abnormal associations in schizophrenia: neural and behavioral evidence. *Neuropsychopharmacology* 33:473–9.
- Juarez I, De La Cruz F, Zamudio S, Flores G. (2005). Cesarean plus anoxia at birth induces hyperresponsiveness to locomotor activity by dopamine D2 agonist. *Synapse* 58: 236–42.
- Juarez I, Silva-Gomez AB, Peralta F, Flores G. (2003). Anoxia at birth induced hyperresponsiveness to amphetamine and stress in postpubertal rats. *Brain Res* 992:281–7.
- Kapur S. (2003). Psychosis as a state of aberrant salience: a framework linking biology, phenomenology, and pharmacology in schizophrenia. *Am J Psychiatry* 160:13–23.
- Katan MB. (1986). Apolipoprotein E isoforms, serum cholesterol, and cancer. *Lancet* 1:507–8.
- Kendler KS, Eaves LJ. (1986). Models for the joint effect of genotype and environment on liability to psychiatric illness. *Am J Psychiatry* 143:279–89.
- Kendler KS, Hewitt J. (1992). The structure of self-report schizotypy in twins. *J Personal Disord* 6:1–17.
- Kendler KS. (2005). “A gene for...”: the nature of gene action in psychiatric disorders. *Am J Psychiatry* 162:1243–52.
- Kendler KS. (2006). Reflections on the relationship between psychiatric genetics and psychiatric nosology. *Am J Psychiatry* 163:1138–46.
- Khoury MJ, Beaty TH, Cohen BH. (1993). *Genetic Epidemiology*. Oxford: Oxford University Press.
- Khoury MJ, Millikan R, Little J, Gwinn M. (2004). The emergence of epidemiology in the genomics age. *Int J Epidemiol* 33:936–44.
- Kirkbride JB, Fearon P, Morgan C, Dazzan P, Morgan K, Tarrant J, Lloyd T, Holloway J, Hutchinson G, Leff JP, Mallett RM, Harrison GL, Murray RM, Jones PB. (2006). Heterogeneity in incidence rates of schizophrenia and other psychotic syndromes: findings from the 3-center AeSOP study. *Arch Gen Psychiatry* 63:250–8.
- Kirkbride JB, Morgan C, Fearon P, Dazzan P, Murray RM, Jones PB. (2007). Neighbourhood-level effects on psychoses: re-examining the role of context. *Psychol Med* 37:1413–25.
- Krabbedam L, van Os J. (2005). Schizophrenia and urbanicity: a major environmental influence – conditional on genetic risk. *Schizophr Bull* 31:795–9.
- Kringle E, Cramer G. (1989). Offspring of monozygotic twins discordant for schizophrenia. *Arch Gen Psychiatry* 46:873–7.
- Kunugi H, Nanko S, Takei N, Saito K, Murray RM, Hirose T. (1996). Perinatal complications and schizophrenia. Data from the Maternal and Child Health Handbook in Japan. *J Nerv Ment Dis* 184:542–6.
- Lake SL, Lyon H, Tantisira K, Silverman EK, Weiss ST, Laird NM, Schaid DJ. (2003). Estimation and tests of haplotype–environment interaction when linkage phase is ambiguous. *Hum Hered* 55:56–65.
- Lange C, DeMeo D, Silverman EK, Weiss ST, Laird NM. (2004). PBAT: tools for family-based association studies. *Am J Hum Genet* 74:367–9.
- Li N, Stephens M. (2003). Modeling linkage disequilibrium and identifying recombination hotspots using single-nucleotide polymorphism data. *Genetics* 165:2213–33.
- Linney YM, Murray RM, Peters ER, MacDonald AM, Rijdsdijk F, Sham PC. (2003). A quantitative genetic analysis of schizotypal personality traits. *Psychol Med* 33:803–16.
- MacDonald AW, Pogue-Geile MF, Debski TT, Manuck S. (2001). Genetic and environmental influences on schizotypy: a community-based twin study. *Schizophr Bull* 27:47–58.
- Malaspina D, Harlap S, Fennig S, Heiman D, Nahon D, Feldman D, Susser ES. (2001). Advancing paternal age and the risk of schizophrenia. *Arch Gen Psychiatry* 58: 361–7.
- Marcelis M, Suckling J, Woodruff P, Hofman P, Bullmore E, van Os J. (2003). Searching for a structural endophenotype in psychosis using computational morphometry. *Psychiatry Res* 122:153–67.

- Marcelis M, Van Os J, Sham P, Jones P, Gilvarry C, Cannon M, McKenzie K, Murray R. (1998). Obstetric complications and familial morbid risk of psychiatric disorders. *Am J Med Genet* 81:29–36.
- McGrath J, Saha S, Welham J, El Saadi O, MacCauley C, Chant D. (2004). A systematic review of the incidence of schizophrenia: the distribution of rates and the influence of sex, urbanicity, migrant status and methodology. *BMC Med* 2:13.
- Meaney MJ, Szyf M. (2005). Environmental programming of stress responses through DNA methylation: life at the interface between a dynamic environment and a fixed genome. *Dialogues Clin Neurosci* 7:103–23.
- Meyer U, Nyffeler M, Schwendener S, Knuesel I, Yee BK, Feldon J. (2008). Relative prenatal and postnatal maternal contributions to schizophrenia-related neurochemical dysfunction after in utero immune challenge. *Neuropsychopharmacology* 33:441–56.
- Meyer-Lindenberg A, Kohn PD, Kolachana B, Kippenhan S, McInerney-Leo A, Nussbaum R, Weinberger DR, Berman KF. (2005). Midbrain dopamine and prefrontal function in humans: interaction and modulation by COMT genotype. *Nat Neurosci* 8:594–6.
- Meyer-Lindenberg A, Miletich RS, Kohn PD, Esposito G, Carson RE, Quarantelli M, Weinberger DR, Berman KF. (2002). Reduced prefrontal activity predicts exaggerated striatal dopaminergic function in schizophrenia. *Nat Neurosci* 5:267–71.
- Meyer-Lindenberg A, Nichols T, Callicott JH, Ding J, Kolachana B, Buckholtz J, Mattay VS, Egan M, Weinberger DR. (2006). Impact of complex genetic variation in COMT on human brain function. *Mol Psychiatry* 11:867–77, 797.
- Moffitt TE, Caspi A, Rutter M. (2005). Strategy for investigating interactions between measured genes and measured environments. *Arch Gen Psychiatry* 62:473–81.
- Mortensen PB, Pedersen CB, Westergaard T, Wohlfahrt J, Ewald H, Mors O, Andersen PK, Melbye M. (1999). Effects of family history and place and season of birth on the risk of schizophrenia [see comments]. *N Engl J Med* 340:603–8.
- Motulsky AG. (1977). Ecogenetics: genetic variation in susceptibility to environmental agents. In: Armendares S, Lisker R. (eds.), *Human Genetics*. Amsterdam, Excerpta Medica, pp. 375–85.
- Murray GK, Corlett PR, Clark L, Pessiglione M, Blackwell AD, Honey G, Jones PB, Bullmore ET, Robbins TW, Fletcher PC. (2008). How dopamine dysregulation leads to psychotic symptoms? Abnormal mesolimbic and mesostriatal prediction error signalling in psychosis. *Mol Psychiatry* 13:239.
- Myin-Germeys I, Marcelis M, Krabbendam L, Delespaul P, van Os J. (2005). Subtle fluctuations in psychotic phenomena as functional states of abnormal dopamine reactivity in individuals at risk. *Biol Psychiatry* 58:105–10.
- Myin-Germeys I, Van Os J, Schwartz JE, Stone AA, Delespaul PA. (2001). Emotional reactivity to daily life stress in psychosis. *Arch Gen Psychiatry* 58:1137–44.
- Nicodemus KK, Marengo S, Batten AJ, Vakkalanka R, Egan MF, Straub RE, Weinberger DR. (2008). Serious obstetric complications interact with hypoxia-regulated/vascular-expression genes to influence schizophrenia risk. *Mol Psychiatry* 13:873–7.
- Nikulina EM, Covington HE, 3rd, Ganschow L, Hammer RP, Jr, Miczek KA. (2004). Long-term behavioral and neuronal cross-sensitization to amphetamine induced by repeated brief social defeat stress: Fos in the ventral tegmental area and amygdala. *Neuroscience* 123:857–65.
- Nimgaonkar VL, Wessely S, Murray RM. (1988). Prevalence of familiarity, obstetric complications, and structural brain damage in schizophrenic patients. *Br J Psychiatry* 153:191–7.
- Norton N, Williams HJ, Owen MJ. (2006). An update on the genetics of schizophrenia. *Curr Opin Psychiatry* 19:158–64.
- O’Callaghan E, Gibson T, Colohan HA, Buckley P, Walshe DG, Larkin C, Waddington JL. (1992). Risk of schizophrenia in adults born after obstetric complications and their association with early onset of illness: a controlled study [see comments]. *BMJ* 305:1256–9.
- O’Donovan MC, Craddock N, Owen MJ. (2008). Schizophrenia: complex genetics, not fairy tales. *Psychol Med* 1–3.

- Ottman R. (1990). An epidemiologic approach to gene–environment interaction. *Genet Epidemiol* 7:177–85.
- Ottman R. (1996). Gene–environment interaction: definitions and study designs. *Prev Med* 25: 764–70.
- Ozawa K, Hashimoto K, Kishimoto T, Shimizu E, Ishikura H, Iyo M. (2006). Immune activation during pregnancy in mice leads to dopaminergic hyperfunction and cognitive impairment in the offspring: a neurodevelopmental animal model of schizophrenia. *Biol Psychiatry* 59:546–54.
- Pedersen CB, Mortensen PB. (2001). Evidence of a dose–response relationship between urbanicity during upbringing and schizophrenia risk. *Arch Gen Psychiatry* 58:1039–46.
- Pedersen CB, Mortensen PB. (2006). Are the cause(s) responsible for urban–rural differences in schizophrenia risk rooted in families or in individuals? *Am J Epidemiol* 163:971–8.
- Plomin R, DeFries JC, Loehlin JC. (1977). Genotype–environment interaction and correlation in the analysis of human behavior. *Psychol Bull* 84:309–22.
- Pulver AE, Liang KY, Brown CH, Wolyniec P, McGrath J, Adler L, Tam D, Carpenter WT, Childs B. (1992). Risk factors in schizophrenia. Season of birth, gender, and familial risk [see comments]. *Br J Psychiatry* 160:65–71.
- Read J, van Os J, Morrison AP, Ross CA. (2005). Childhood trauma, psychosis and schizophrenia: a literature review with theoretical and clinical implications. *Acta Psychiatr Scand* 112:330–50.
- Rutter M, Moffitt TE, Caspi A. (2006). Gene–environment interplay and psychopathology: multiple varieties but real effects. *J Child Psychol Psychiatry* 47:226–61.
- Rutter M. (2005). How the environment affects mental health. *Br J Psychiatry* 186:4–6.
- Rutter M. (2006). *Genes and Behaviour: Nature–Nurture Interplay Explained*. Malden, MA, USA: Blackwell Publishing.
- Schmidt-Kastner R, van Os J, Steinbusch HWMS, Schmitz C. (2006). Gene regulation by hypoxia and the neurodevelopmental origin of schizophrenia. *Schizophr Res* 84:253–71.
- Schurhoff F, Szoke A, Chevalier F, Roy I, Meary A, Bellivier F, Giros B, Leboyer M. (2007). Schizotypal dimensions: an intermediate phenotype associated with the COMT high activity allele. *Am J Med Genet B Neuropsychiatr Genet* 144:64–8.
- Schurhoff F, Szoke A, Meary A, Bellivier F, Rouillon F, Pauls D, Leboyer M. (2003). Familial aggregation of delusional proneness in schizophrenia and bipolar pedigrees. *Am J Psychiatry* 160:1313–9.
- Selten JP, Cantor-Graae E, Kahn RS. (2007). Migration and schizophrenia. *Curr Opin Psychiatry* 20:111–5.
- Sham P. (1996). Genetic epidemiology. *Br Med Bull* 52:408–33.
- Shur E. (1982). Season of birth in high and low genetic risk schizophrenics. *Br J Psychiatry* 140:410–5.
- Sipos A, Rasmussen F, Harrison G, Tynelius P, Lewis G, Leon DA, Gunnell D. (2004). Paternal age and schizophrenia: a population based cohort study. *BMJ* 329:1070.
- Smits L, Pedersen C, Mortensen P, Van Os J. (2004). Association between short birth intervals and schizophrenia in the offspring. *Schizophr Res* 70:49–56.
- Spauwen J, Krabbendam L, Lieb R, Wittchen HU, Van Os J. (2006a). Evidence that the outcome of developmental expression of psychosis is worse for adolescents growing up in an urban environment. *Psychol Med* 407–15.
- Spauwen J, Krabbendam L, Lieb R, Wittchen HU, van Os J. (2006b). Impact of psychological trauma on the development of psychotic symptoms: relationship with psychosis proneness. *Br J Psychiatry* 188:527–33.
- Stefanis NC, Henquet C, Avramopoulos D, Smyrnis N, Evdokimidis I, Myin-Germeyns I, Stefanis CN, Van Os J. (2007). COMT Val158Met moderation of stress-induced psychosis. *Psychol Med* 37:1651–6.
- Stefanis NC, Van Os J, Avramopoulos D, Smyrnis N, Evdokimidis I, Hantoumi I, Stefanis CN. (2004). Variation in catechol-o-methyltransferase val158 met genotype associated with schizotypy but not cognition: a population study in 543 young men. *Biol Psychiatry* 56: 510–5.

- Sugarman PA, Craufurd D. (1994). Schizophrenia in the Afro-Caribbean community. *Br J Psychiatry* 164:474–80.
- Sullivan PF. (2008). The dice are rolling for schizophrenia genetics. *Psychol Med* 1–4.
- Susser E, Neugebauer R, Hoek HW, Brown AS, Lin S, Labovitz D, Gorman JM. (1996). Schizophrenia after prenatal famine. Further evidence [see comments]. *Arch Gen Psychiatry* 53:25–31.
- Susser E, Susser M. (1989). Familial aggregation studies. A note on their epidemiologic properties. *Am J Epidemiol* 129:23–30.
- Talkowski ME, Kirov G, Bamne M, Georgieva L, Torres G, Mansour H, Chowdari KV, Milanova V, Wood J, McClain L, Prasad K, Shirts B, Zhang J, O'Donovan MC, Owen MJ, Devlin B, Nimgaonkar VL. (2008). A network of dopaminergic gene variations implicated as risk factors for schizophrenia. *Hum Mol Genet* 17:747–58.
- Tenn CC, Fletcher PJ, Kapur S. (2005). A putative animal model of the “prodromal” state of schizophrenia. *Biol Psychiatry* 57:586–93.
- Tidey JW, Miczek KA. (1996). Social defeat stress selectively alters medocorticolimbic release: an in vivo microdialysis study. *Brain Research* 721:140–9.
- Tienari P, Wynne LC, Moring J, Lahti I, Naarala M, Sorri A, Wahlberg KE, Saarento O, Seitamaa M, Kaleva M, et al. (1994). The Finnish adoptive family study of schizophrenia. Implications for family research [see comments]. *Br J Psychiatry Suppl*:20–6.
- Tienari P, Wynne LC, Sorri A, Lahti I, Laksy K, Moring J, Naarala M, Nieminen P, Wahlberg KE. (2004). Genotype–environment interaction in schizophrenia-spectrum disorder. Long-term follow-up study of Finnish adoptees. *Br J Psychiatry* 184:216–22.
- Toulopoulou T, Picchioni M, Rijdsdijk F, Hua-Hall M, Ettinger U, Sham P, Murray R. (2007). Substantial genetic overlap between neurocognition and schizophrenia: genetic modeling in twin samples. *Arch Gen Psychiatry* 64:1348–55.
- van Haren NE, Bakker SC, Kahn RS. (2008). Genes and structural brain imaging in schizophrenia. *Curr Opin Psychiatry* 21:161–7.
- Van Os J, Fahy TA, Bebbington P, Jones P, Wilkins S, Sham P, Russell A, Gilvarry K, Lewis S, Toone B, et al. (1994). The influence of life events on the subsequent course of psychotic illness. A prospective follow-up of the Camberwell Collaborative Psychosis Study. *Psychol Med* 24:503–13.
- Van Os J, Hanssen M, Bak M, Bijl RV, Vollebergh W. (2003). Do urbanicity and familial liability coparticipate in causing psychosis? *Am J Psychiatry* 160:477–82.
- Van Os J, Hanssen M, Bijl R-V, Ravelli A. (2000). Straus (1969) revisited: a psychosis continuum in the general population? *Schizophr Res* 45:11–20.
- van Os J, Henquet C, Stefanis N. (2005). Cannabis-related psychosis and the gene–environment interaction: comments on Ferdinand et al. 2005. *Addiction* 100:874–5.
- Van Os J, Krabbendam L, Myin-Germeys I, Delespaul P. (2005). The schizophrenia envirome. *Curr Opin Psychiatry* 18:141–5.
- Van Os J, Marcelis M. (1998). The ecogenetics of schizophrenia: a review. *Schizophr Res* 32:127–35.
- van Os J, Pedersen CB, Mortensen PB. (2004). Confirmation of synergy between urbanicity and familial liability in the causation of psychosis. *Am J Psychiatry* 161:2312–4.
- Van Os J, Sham P. (2003). Gene–environment interactions. In: Murray RM, Jones PB, Susser E, Van Os J, Cannon M. (eds.), *The Epidemiology of Schizophrenia*. Cambridge, Cambridge University Press, pp. 235–54.
- Van Os J. (2004). Does the urban environment cause psychosis? *Br J Psychiatry* 184:287–8.
- van Os J. (2005). Commentary on residential location papers by Whitfield et al. (2005) and Willemssen et al. (2005). *Twin Res Hum Genet* 8:318–9.
- Venerosi A, Valanzano A, Cirulli F, Alleva E, Calamandrei G. (2004). Acute global anoxia during C-section birth affects dopamine-mediated behavioural responses and reactivity to stress. *Behav Brain Res* 154:155–64.

- Verdoux H, Gindre C, Sorbara F, Tournier M, Swendsen JD. (2003). Effects of cannabis and psychosis vulnerability in daily life: an experience sampling test study. *Psychol Med* 33:23–32.
- Verdoux H, Van Os J, Sham P, Jones P, Gilvarry K, Murray R. (1996). Does familiarity predispose to both emergence and persistence of psychosis? A follow-up study. *Br J Psychiatry* 168:620–6.
- Vollema MG, Sitskoorn MM, Appels MC, Kahn RS. (2002). Does the Schizotypal Personality Questionnaire reflect the biological-genetic vulnerability to schizophrenia? *Schizophr Res* 54:39–45.
- Wahlberg KE, Wynne LC, Hakko H, Laksy K, Moring J, Miettunen J, Tienari P. (2004). Interaction of genetic risk and adoptive parent communication deviance: longitudinal prediction of adoptee psychiatric disorders. *Psychol Med* 34:1531–41.
- Wahlberg KE, Wynne LC, Oja H, Keskitalo P, Pykalainen L, Lahti I, Moring J, Naarala M, Sorri A, Seitamaa M, Laksy K, Kolassa J, Tienari P. (1997). Gene–environment interaction in vulnerability to schizophrenia: findings from the Finnish Adoptive Family Study of Schizophrenia. *Am J Psychiatry* 154:355–62.
- Walker E. (1981). Attentional and neuromotor functions of schizophrenics, schizoaffectives, and patients with other affective disorders. *Arch Gen Psychiatry* 38:1355–8.
- Weaver IC, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, Dymov S, Szyf M, Meaney MJ. (2004). Epigenetic programming by maternal behavior. *Nat Neurosci* 7:847–54.
- Weiser M, van Os J, Reichenberg A, Rabinowitz J, Nahon D, Kravitz E, Lubin G, Shmushkevitz M, Knobler HY, Noy S, Davidson M. (2007). Social and cognitive functioning, urbanicity and risk for schizophrenia. *Br J Psychiatry* 191:320–4.
- Weisfeld GE, Weisfeld CC. (2002). Marriage: an evolutionary perspective. *Neuro Endocrinol Lett* 23(Suppl 4):47–54.
- Wellman CL, Izquierdo A, Garrett JE, Martin KP, Carroll J, Millstein R, Lesch KP, Murphy DL, Holmes A. (2007). Impaired stress-coping and fear extinction and abnormal corticolimbic morphology in serotonin transporter knock-out mice. *J Neurosci* 27:684–91.
- Whitfield JB, Zhu G, Heath AC, Martin NG. (2005). Choice of residential location: chance, family influences, or genes? *Twin Res Hum Genet* 8:22–6.
- Wichers M, Myin-Germeys I, Jacobs N, Peeters F, Kenis G, Derom C, Vlietinck R, Delespaul P, Van Os J. (2007a). Genetic risk of depression and stress-induced negative affect in daily life. *Br J Psychiatry* 191:218–23.
- Wichers MC, Myin-Germeys I, Jacobs N, Peeters F, Kenis G, Derom C, Vlietinck R, Delespaul P, van Os J. (2007b). Evidence that moment-to-moment variation in positive emotions buffer genetic risk for depression: a momentary assessment twin study. *Acta Psychiatr Scand* 115:451–7.
- Wilkinson LS, Davies W, Isles AR. (2007). Genomic imprinting effects on brain development and function. *Nat Rev Neurosci* 8:832–43.
- Willemsen G, Posthuma D, Boomsma DI. (2005). Environmental factors determine where the Dutch live: results from the Netherlands twin register. *Twin Res Hum Genet* 8:312–7.
- Wong MY, Day NE, Luan JA, Chan KP, Wareham NJ. (2003). The detection of gene–environment interaction for continuous traits: should we deal with measurement error by bigger studies or better measurement? *Int J Epidemiol* 32:51–7.
- Yui K, Goto K, Ikemoto S, Ishiguro T. (2000). Stress induced spontaneous recurrence of methamphetamine psychosis: the relation between stressful experiences and sensitivity to stress. *Drug Alcohol Depend* 58:67–75.
- Zammit S, Allebeck P, Dalman C, Lundberg I, Hemmingson T, Owen MJ, Lewis G. (2003). Paternal age and risk for schizophrenia. *Br J Psychiatry* 183:405–8.
- Zammit S, Lewis S, Gunnell D, Smith GD. (2007). Schizophrenia and neural tube defects: comparisons from an epidemiological perspective. *Schizophr Bull* 33: 853–8.
- Zinkstok J, van Nimwegen L, van Amelsvoort T, de Haan L, Yusuf MA, Baas F, Linszen D. (2008). Catechol-*O*-methyltransferase gene and obsessive-compulsive symptoms in patients with recent-onset schizophrenia: preliminary results. *Psychiatry Res* 157:1–8.